arXiv:cond-mat/0612063v2 [cond-mat.dis-nn] 17 Dec 2006

An invariant distribution in static granular media

T. Aste, T. Di Matteo, M. Saadatfar, T.J. Senden
Department of Applied Mathematics, The Australian National University, 0200 Canberra, ACT, Australia

M. Schröter and H. L. Swinney
Center for Nonlinear Dynamics and Department of Physics,
University of Texas at Austin, Austin, Texas 78712 USA

We have discovered an invariant distribution for local packing configurations in static granular media. This distribution holds in experiments for packing fractions covering most of the range from random loose packed to random close packed, for beads packed both in air and in water. Assuming only that there exist elementary cells in which the system volume is subdivided, we derive from statistical mechanics a distribution that is in accord with the observations. This universal distribution function for granular media is analogous to the Maxwell-Boltzmann distribution for molecular gasses.

PACS numbers: 45.70.-n Granular Systems 45.70.Cc Static sandpiles; Granular Compaction 81.05.Rm-Porous materials; granular materials

Granular materials are complex systems characterized by unusual static and dynamic properties. These systems are comprised of large numbers of dissipative macroscopic particles assembled into disordered structures. The microscopic description of the system-state requires a very large number of variables. However, there is a very large number of microscopic configurations corresponding to the same macroscopic properties. Edwards and coauthors [1, 2] have proposed that the complexity of static granular systems could be disentangled by means of a statistical mechanics approach reducing the description of the system state to a few parameters only [3, 4, 5, 6, 7, 8, 9, 10, 11, 12]. An essential part of Edwards' idea is that in static granular media volume plays the role held by energy in usual thermodynamics. Therefore, an understanding of the volume distribution function is the key to connect microscopic details of the system with macroscopic state variables.

Since granular materials are dissipative, they can change their static configurations only when energy is injected into the system. The general idea underlying a statistical mechanics description is that the properties of the system do not depend on the kind of energy injections, but only on the portion of the configurational space that the system explores under such action. For instance, some classical experiments [13, 14, 15] obtained different average packing fractions by tapping the container with different intensities and different numbers of times. Similarly, in an experiment by Schröter et al. [16], reproducible average packing fractions were obtained by driving the system with periodic trains of flow pulses in a fluidized bed. More generally, various controlled perturbations (tapping, rotating, pouring, etc.) can produce packings with characteristic average packing fractions. Within a statistical mechanics framework, if the system volume (V_T) is the relevant state variable, the system properties should depend only on the achieved packing fraction and not on the preparation history. This hypothesis is examined in this paper using the largest sets of experimental data on particle positions presently available.

Experiments.- We analyze the structural properties of static granular packings produced in 18 different experiments, 6 with acrylic spheres in air and 12 with glass beads in water. The packing fractions ρ range from 0.56 to 0.64. Three-dimensional density maps have been obtained for these systems using X-ray Computed Tomography [17]. Coordinates of the bead centers have been calculated for more than two million spheres with a precision better than 1% of their diameters, which is better than the uncertainty arising from polydispersity (5% for the glass particles and 2% for the acrylic particles).

The 12 samples of glass beads were prepared using the fluidized bed technique described in [16]; each sample consisted of about 145,000 spherical grains with diameters $250 \pm 13\mu$m placed in a cylindrical glass container with inner diameter 12.7 mm. The beads were fluidized with pulses of deionized water with the flow rate pulsed by a computer-controlled syringe pump. During each flow pulse the bed expanded until its height reached a stable value. After each flow pulse the bed was allowed to settle into a mechanically stable configuration. Packing fractions in the range between 0.56 and 0.60 were obtained by tuning the water flux. The particle configurations have been studied for sub-sets of grains that were at a distance larger than four sphere diameters from the sample boundaries; there were about 90,000 grains per sample.

The 6 samples with acrylic spheres were described in [18]. In particular, two samples contained 140,000 particles (90,000 in the internal volume) with diameter 1.59 ± 0.02 mm in a cylindrical container with an inner diameter of 55 mm, filled to a height of 75 mm. The other four samples each had about 35,000 (15,000 in the internal volume) acrylic spheres with diameter 1.00 ± 0.02 mm in the same cylindrical container.

Theory.- Since the original work by Edwards and coauthors, several statistical mechanics approaches have



FIG. 1: (a) Distributions of the Delaunay cell volumes from 18 experiments collapse onto a universal curve when plotted vs. $(V - V_{min})/(\langle V \rangle - V_{min})$. The inset shows the same distributions plotted vs. V/d^3. (b) The inverse cumulate distributions, $F(V) = 1 - \int_{V_{min}}^{V} f(V)dV$, also show evidence of a collapse onto a universal curve when plotted vs. $(V - V_{min})/(\langle V \rangle - V_{min})$. The line is the inverse cumulate distributions for $f(V, k = 1)$ (Eq.1). The inset shows the cumulate distributions plotted versus V/d^3. The distributions are obtained from a statistical analysis over more than six million cells in 18 different experiments with three different kinds of beads in air and in water. The open symbols correspond to the 6 experiments with with dry acrylic beads [18]: circles sample A ($\rho = 0.586$); squares sample B ($\rho = 0.596$); stars sample C ($\rho = 0.619$); diamonds sample D ($\rho = 0.626$); triangles up sample E ($\rho = 0.630$); triangles down sample F ($\rho = 0.64$). The '+' corresponds to the 12 experiments with packing fractions $0.56 \leq \rho \leq 0.60$ made with glass spheres in of fluidized bed.

been proposed to describe granular materials [1, 2, 4, 5, 6, 7, 8, 10, 19]. We make here the minimal number of assumptions in considering a granular system at rest that occupies a given volume V_T. We assume that the system can be subdivided into C elementary cells with volumes v_i $(i = 1...C)$ with $\sum_{i=1}^{C} v_i = V_T$. Under the sole assumption that any assembly of such elementary volumes will produce proper mechanically stable packings, we obtain the probability distribution for the cell-elementary volumes:

$$p(v) = \frac{1}{\chi} e^{-(v - v_{min})/\chi} \quad , \tag{1}$$

with

$$\chi = \langle v \rangle - v_{min} \quad , \tag{2}$$

where $\langle v \rangle = V_T/C$ is the average volume per elementary cell, v_{min} is the minimum volume attainable by the elementary cells and χ is an intensive thermodynamic parameter accounting for the exchange of volume between the elementary cell and the surrounding volume "reservoir". There is no need to specify the nature and kind of such "elementary cells"; the only assumptions are that they exist, that they can have any volume larger than v_{min}, and that their assembly fills a volume V_T. Space can be divided arbitrarily into pieces. Common examples of such partitions into space-filling blocks are the Delaunay and the Voronoï decompositions [20, 21, 22]. Such cells do not coincide with the elementary ones, but they might be assemblies of such elementary cells. For instance, in a Delaunay partition of a three-dimensional packing there are about six times more cells than in the Voronoï decomposition. The present theory applies to any degree of space partition made by any agglomerate local structure made of a given number k of elementary cells. From Eq. 1 the probability distribution for the volumes of such agglomerate of k elementary cells is

$$f(V, k) = \frac{k^k}{(k-1)!} \frac{(V - V_{min})^{(k-1)}}{(\langle V \rangle - V_{min})^k} \exp\left(-k \frac{V - V_{min}}{\langle V \rangle - V_{min}}\right) \quad , \tag{3}$$

with $\langle V \rangle = V_T/C$ and $V_{min} = k v_{min}$ being respectively the average and the minimum volumes for a given packing. It follows from Eqs. 1 and 3 that

$$\chi = \frac{\langle V \rangle - V_{min}}{k} \quad , \tag{4}$$

which is the average free-volume per elementary cell. Therefore, the intensive variable χ is a measure of the kind and the degree of space-partition into elementary cells.

We have obtained $f(V, k)$ assuming that the cells are uniquely characterized by their volumes, and that any combination of C cells with arbitrary volumes $v_i \geq v_{min}$ will produce a structurally stable, space-filling system of cells. This is possible in one-dimension only where, indeed, the packing is an arbitrary disposition of grain centers at distances larger or equal than v_{min}. In this case, the elementary cells are the Delaunay cells [22], which are the segments in between two successive grain centers; the distribution of their sizes is exactly described by $p(v)$. In three dimensions, the Delaunay cells are tetrahedra with vertices on the centers of neighborhood grains chosen in a way that no other grains in the packing have centers within the circumsphere of each Delaunay tetrahedron. Clearly, in this case, the Delaunay cells are not uniquely described by their volumes and an arbitrary collection of cells is neither space-filling nor mechanically stable.

Results.- We observe that the distributions of the Delaunay volumes obtained from the 18 sets of data for



FIG. 2: The distributions of the Voronoï cell volumes collapse onto a universal curve when plotted vs. $(V-V_{min})/(\langle V\rangle-V_{min})$. The theoretical line is $f(V,k=12)$ (Eq. 1). The inset shows the distributions plotted versus V/d^3. The data and the symbols are the same as in Fig. 1.

different kinds of beads in different media and different conditions are all described well by the *same* function, as shown Fig.1(a). The collapse of the data onto a single curve was obtained using $V_{min}=\sqrt{2}/12d^3$ (with d the sphere diameter), which is the volume of a regular Delaunay tetrahedron for four spheres in contact (the smallest compact tetrahedron [21]). The tail of the observed distribution function for the Delauney volumes is exponential, as in Eq. 1 with $k\sim 1$. The data fit well to the inverse cumulate distribution, $F(V)=1-\int_{v_{min}}^{V}f(v',1)dv'$, as Fig.1(b) shows; the agreement extends over four orders of magnitude and uses no adjustable parameters. The agreement suggests that Delaunay cells might be considered candidates for the "elementary cells"; however, it was noted in [21] that, at small $V-V_{min}$ in the region where the spheres make contact ($V<\sqrt{3}d^3/12$), the empirical distributions deviate from the simple exponential form predicted by $p(v)$, which indicates that other constraints such as mechanical equilibrium should be also taken into account to correctly describe such region.

An alternative way for dividing space into space-filling cells is the Voronoï partition. In one dimension, the Voronoï cell around grain 'i' is constructed by taking the segment between the two mid-points between grains $i-1$ and i, and between grains i and $i+1$. The size of such a segment is equal to $(v_i+v_{i+1})/2$ where v_i and v_{i+1} are the distances between the two consecutive couples of points. Therefore, the probability of finding a one-dimensional Voronoï cell of size V is associated with the probability of finding two successive Delaunay cells with sizes $v_1+v_2=2V$. This probability is $\int_{2v_{min}}^{V-v_{min}}p(v_1)p(v-v_1)dv_1\propto(V-2v_{min})\exp[-(V-2v_{min})/(\langle V\rangle-2v_{min})]\propto f(V,k=2)$, with $\langle V\rangle$ the average size of the Voronoï cell. In three dimensions, the Voronoï cell can be also seen as the combination of several elementary cells with distribution $f(V,k)$; however, in this case the number of sub-cells involved is not fixed at $k=2$ but depends on the kind of packing.



FIG. 3: The standard deviations of the Voronoï volume distributions in the 18 experiments fit the linear relation given by Eq. 6 with $k=12$. The dash line above (below) the solid line corresponds to $k=11$ ($k=13$).

We observe that data for over a million of Voronoï cells from all 18 experiments for dry and wet packings of glass and acrylic spheres collapse to the the same distribution function (see Fig. 2). In this collapse of the data there no adjustable paramaters, just $\langle V\rangle=V_T/$ (*number of grains*) and $V_{min}=5^{(5/4)}/\sqrt{2(29+13\sqrt{5})}d^3\simeq 0.694d^3$, which is the smallest Voronoï cell that can be built in a equal-spheres packing [22]. Figure 2 shows that such universal distribution function is well described by Eq. 1 with $k=12$, which indicates that about 12 elementary cells contribute in building each Voronoï cell. Such a number is meaningful, since about 12 spheres are expected to be found in the close neighboring of any given sphere in the packing.

A further demonstration that such statistical distributions are independent of the details of a sample and the method of sample preparation is provided by the behavior of the volume fluctuations, which can be calculated directly from the relation (see Eq. 1),

$$\sigma^2(V)=\left\langle(V-\langle V\rangle)^2\right\rangle=\chi^2\frac{\partial\langle V\rangle}{\partial\chi}\quad , \tag{5}$$

which becomes, using $\partial\langle V\rangle/\partial\chi=k$ (from Eq. 4),

$$\sigma(V)=\sqrt{k}\ \chi=\frac{\langle V\rangle-V_{min}}{\sqrt{k}}\quad . \tag{6}$$

The good correspondence of the data from the Voronoï volume distributions with Eq. 6 (Fig. 3) provides further evidence that $\langle V\rangle$ and k are the relevant control parameters, independent of the system and its preparation.

Conclusion.- We have shown that for a broad set of granular packings of different monodisperse spherical grains, prepared by using different procedures, the local volumes are described by a universal distribution function (Eq. 3), which can be predicted from a minimal set of assumptions. The only tunable parameter in the present theory is k, which has the value 12 for Voronoï decompositions throughout the accessible density ranges of the different static granular packings studied. Interestingly, for granular gasses the same empirical distribution (Eq. 3) applies but with $k = 5.586$ [23, 24]. A similar kind of distribution (Gamma distribution) with different values of the parameter k have also been observed in two dimensional Voronoï tessellation generated from disk packings [25]. Therefore, the parameter k could be the 'structure parameter' which depends on the system-phase. Within the framework of a statistical mechanics description, Eq. 3 can be regarded as the analogous to the Maxwell-Boltzmann distribution for granular media.

We thank Mario Nicodemi, Massimo Pica Ciamarra, Tom Truskett and Philip Metzger for helpful discussions and comments. This work was partially supported by the ARC discovery project DP0450292. MS and HLS were supported by the Robert A. Welch Foundation.

---

[1] S. Edwards and R. Oakeshott, Physica A **157**, 1080 (1989).

[2] A. Mehta and S. F. Edwards, Physica A **157**, 1091 (1989).

[3] A. Fierro, M. Nicodemi, and A. Coniglio, Europhys. Lett. **59**, 642 (2002).

[4] H. A. Makse and J. Kurchan, Nature **415**, 614 (2002).

[5] B. Behringer, Nature **415**, 594 (2002).

[6] G. D'Anna, P. Mayor, A. Barrat, V. Loreto, and F. Nori, Nature **424**, 909 (2003).

[7] R. P. Ojha, P. A. Lemieux, P. K. Dixon, A. J. Liu, and D. J. Durian, Nature **427**, 521 (2004).

[8] P. Richard, M. Nicodemi, R. Delannay, P. Ribiere, and D. Bideau, Nature Materials **4**, 121 (2005).

[9] M. P. Ciamarra, M. Nicodemi, and A. Coniglio, Phys. Rev. Lett. **97**,158001 (2006).

[10] F. Lechenault, F. da Cruz, O. Dauchot, and E. Bertin, J. Stat. Mech. **P07009**, 1742 (2006).

[11] P. T. Metzger, Phys. Rev. E. **70**,051303 (2004).

[12] P. T. Metzger and C. M. Donahue, Phys. Rev. Lett. **94**, 148001 (2005).

[13] J. B. Knight, C. G. Fandrich, C. N. Lau, H. M. Jeager, and S. R. Nagel, Phys Rev E **51**, 3957 (1995).

[14] E. R. Nowak, J. B. Knight, M. L. Povinelli, H. M. Jeager, and S. R. Nagel, Powder Technol. **94**, 79 (1997).

[15] E. R. Nowak, J. B. Knight, E. BenNaim, H. M. Jeager, and S. R. Nagel, Phys. Rev. E **57**, 1971 (1998).

[16] M. Schröter, D. I. Goldman, and H. L. Swinney, Phys. Rev. E. **71**, 30301 (R) (2005).

[17] T. Aste, M. Saadatfar, A. Sakellariou, and T. Senden, Physica A **339**, 16 (2004).

[18] T. Aste, M. Saadatfar, and T. J. Senden, Phys. Rev. E. **71**, 061302 (2005).

[19] M. P. Ciamarra, M. Nicodemi, and A. Coniglio, preprint. (2006).

[20] G. Voronoï, J. reine angew. Math. **134**, 198 (1908).

[21] T. Aste, Phys. Rev. Lett. **96**, 018002 (2006).

[22] T. Aste and D. Weaire, *The Pursuit of Perfect Packing* (Institute of Physics, Bristol, 2000).

[23] E. N. Gilbert, Ann. Math. Stat. **33**, 958 (1962).

[24] E. Pineda, P. Bruna, and D. Crespo, Phys. Rev. E. **70**, 066119 (2004).

[25] V. S. Kumar and V. Kumaran, J. Chem. Phys. **123**, 114501 (2005).